

Pawitar Dulari

Department of Physics, Government Degree College Dharamshala (H.P.) INDIA-176215
*e-mail:pawitar.ibs@gmail.com

**Thermoelastic properties of solids
based on equation of state**

Abstract. Thermoelastic properties of solids at high pressures are studied using various equations of state (EOS) such as Eularian Birch-Murnaghan EOS, Poirier-Tarantola logarithmic EOS and the generalized Vinet-Rydberg EOS. We have determined the pressure derivatives of bulk modulus upto third order which are useful for predicting the Grüneisen parameter and its volume derivatives. Expressions have been obtained for the derivative properties based on different equations of state, and extrapolated to the limit of extreme compression. It is found that all the three equations lead to a common relationship between second and third pressure derivatives of bulk modulus in the limit of extreme compression.

Keywords: Equations of state, pressure derivatives of bulk modulus, Grüneisen parameter, extreme compression behavior.

Introduction

Equations of state at high pressures have been extremely useful for studying the thermoelastic properties of solids [1-5]. Bulk modulus and its pressure derivatives are important physical quantities for understanding the thermoelastic properties [6, 7] such as the Grüneisen parameter γ and its volume derivatives. γ is related to the thermal and elastic properties of materials by the formula [8, 9].

$$\gamma = \frac{\alpha K_T V}{C_V} = \frac{\alpha K_S V}{C_P} \quad (1)$$

Where α is the coefficient of volume thermal expansion, K_T and K_S are isothermal and adiabatic bulk modulus, C_V and C_P are the specific heats at constant pressure and constant volume respectively.

The second Grüneisen constant q used in the literature is defined as [10].

$$q = \left(\frac{\partial \ln \gamma}{\partial \ln V} \right)_T \quad (2)$$

The third order Grüneisen parameter is defined as

$$\lambda = \left(\frac{\partial \ln q}{\partial \ln V} \right)_T \quad (3)$$

or

$$\lambda = 1 - q + \frac{V}{q\gamma} + \frac{d^2\gamma}{dV^2} \quad (4)$$

In order to emphasize the importance of q and λ in determining higher order thermoelastic properties we refer to the following thermodynamic identities [9].

$$\delta_S = K'_S - 1 + q - \gamma - C'_S \quad (5)$$

$$\delta_T = K'_T - 1 + q + C'_T \quad (6)$$

$$\begin{aligned} & \left(\frac{\partial \delta_S}{\partial \ln V} \right)_S - K_S K''_S + q(\lambda - \gamma) + (\gamma + q) \\ & \left(\frac{\partial \ln C_V}{\partial \ln V} \right)_S - \left(\frac{\partial}{\partial \ln V} \left(\frac{\partial \ln C_V}{\partial \ln V} \right)_S \right)_S - \\ & - \left(\frac{\partial}{\partial \ln V} \left(\frac{\partial \ln C_V}{\partial \ln V} \right)_S \right)_T \end{aligned} \quad (7)$$

$$\left(\frac{\partial \delta_s}{\partial \ln V}\right)_T = -K_T K_T'' + \lambda q + \left(\frac{\partial C_T'}{\partial \ln V}\right)_T \quad (8)$$

where

δ_s is the adiabatic Anderson-Grüneisen parameter

$$\delta_s = -\frac{1}{\alpha} \left(\frac{\partial \ln K_s}{\partial T}\right)_P \quad (9)$$

δ_T is the isothermal Anderson-Grüneisen parameter

$$\delta_T = -\frac{1}{\alpha} \left(\frac{\partial \ln K_T}{\partial T}\right)_P \quad (10)$$

$$C_s' = (\partial \ln C_V / \partial \ln V)_S \quad (11)$$

and

$$C_T' = (\partial \ln C_V / \partial \ln V)_T \quad (12)$$

Thus q and λ appearing in equations [5–8] are useful parameters reduced to investigate higher order thermoelastic properties. We make use of the generalized free-volume formula for determining q and λ .

Generalized free-volume formulation

According to the generalized free-volume formula [8, 11], γ is related to pressure P , isothermal bulk modulus K and its pressure derivative K' as follows:

$$\gamma = \frac{\left(\frac{K'}{2}\right) - \left(\frac{1}{6}\right) - \left(\frac{t}{3}\right) \left(1 - \frac{P}{3K}\right)}{1 - 2t \left(\frac{P}{3K}\right)} \quad (13)$$

The parameter t takes different values for different derivatives of γ , based on different approximation. Thus $t = 0$ for Slater's formula [12], $t = 1$ for the formulation developed by Dugdale and MacDonald [13], $t = 2$ yields the free-volume formula [9], and $t = 2.35$ resulted in a molecular dynamical calculation by Barton and Stacey [14]. The assumptions and approximations on which Equation [13] is based, have been reviewed in a comprehensive manner by Stacey and Davis. Equation [13] can be applied to different types of metals, solids as well as insulators, because it is

derived from the fundamental relationship between thermal pressure and thermal energy [8]. The pressure dependence or volume dependence of γ can be studied with the help of Eq. [13] using different equations of state [10, 15]. Expressions for the volume derivatives of γ , represented by q and λ are derived from Eq. (13) considering t to be independent of pressure, i.e. $dt/dp_T = 0$. It has been found by Stacey and Davis [8] that λ varies slowly with pressure, and constant λ might be a good approximation. Although there is no fundamental reason for believing that λ is constant, it is much better assumption constant q often assumed in mineral physics [16].

It is more convenient to rewrite Eq. (13) in an equivalent form as follows (13).

$$\gamma = \frac{K'}{2} - \frac{1}{6} - \varepsilon \quad (14)$$

where

$$\varepsilon = \frac{t(K - K'P)}{(3K - 2tP)} \quad (15)$$

The following equations are then obtained from the differentiation of Eq. (13)

$$\gamma_q = \frac{-KK''}{2} + K \frac{d\varepsilon}{dP} \quad (16)$$

and

$$\gamma_q (q + \lambda) = \frac{K^2 K'''}{2} + \frac{K'(KK'')}{2} - K' \left(K \frac{d\varepsilon}{dP} \right) - K^2 \frac{d^2 \varepsilon}{dP^2} \quad (17)$$

Eqs. (16) and (17) yield

$$(q + \lambda) = -K' - \frac{[(K^2 K''' / KK'') - (2 / KK'') (K^2 d^2 \varepsilon / dP^2)]}{1 - (2 / KK'') (K d\varepsilon / dP)} \quad (18)$$

Values of γ , q and λ can be calculated by knowing the pressure derivatives of bulk modulus. These pressure derivatives can be determined with the help of equations of state.

Analysis based on equations of state

Higher pressure derivatives of bulk modulus are determined here using some important equations of state given below:

Birch-Murnaghan fourth order EOS

This EOS has been derived from the Eulerian strain theory [17]. The expressions for P, K, K', KK'' and K²K''' obtained from this equation of state are given below:

$$P = \frac{9K_0}{16} [-Ax^{-5/3} + Bx^{-7/3} - Cx^{-3} + Dx^{-11/3}], \quad (19)$$

$$K = \frac{9K_0}{16} [-A(5/3)x^{-5/3} + B(7/3)x^{-7/3} - C(3)x^{-3} + D(11/3)x^{-11/3}] \quad (20)$$

$$K' = \frac{9K_0}{16K} [-A(5/3)^2x^{-5/3} + B(7/3)^2x^{-7/3} - C(3)^2x^{-3} + D(11/3)^2x^{-11/3}] \quad (21)$$

$$KK'' = \frac{9K_0}{16K} [-A(5/3)^3x^{-5/3} + B(7/3)^3x^{-7/3} - C(3)^3x^{-3} + D(11/3)^3x^{-11/3}] - K^{12} \quad (22)$$

$$K^2 K''' = \frac{9K_0}{16K} [-A(5/3)^4x^{-5/3} + B(7/3)^4x^{-7/3} - C(3)^4x^{-3} + D(11/3)^4x^{-11/3}] - K^{13} - 4K'KK'' \quad (23)$$

where $x = V/V_0$ and

$$A = K_0 K_0'' + (K_0' - 4)(K_0' - 5) + 59/9 \quad (24)$$

$$B = 3 K_0 K_0'' + (K_0' - 4)(3K_0' - 13) + 129/9 \quad (25)$$

$$C = 3 K_0 K_0'' + (K_0' - 4)(3K_0' - 11) + 105/9 \quad (26)$$

$$D = K_0 K_0'' + (K_0' - 4)(K_0' - 3) + 35/9 \quad (27)$$

Poirier-Tarantola logarithmic fourth-order EOS

Poirier and Tarantola [18] have obtained logarithmic EOS using the Hencky strain which is represented by $(1/3) (\ln V/V_0)$. The expressions based on this EOS s follows:

$$P = K_0 x^{-1} [-(\ln x) + \left(\frac{K_0' - 2}{2}\right)(\ln x)^2 - \frac{1}{6}Q(\ln x)^3] \quad (28)$$

$$K = K_0 x^{-1} [1 - (K_0' - 1)(\ln x) + \frac{1}{2}(K_0 K_0'' + K_0^{12} + 2K_0' + 1)(\ln x)^2 - \frac{1}{6}Q(\ln x)^3] \quad (29)$$

$$K' = \frac{K_0 x^{-1}}{K} [K_0' - (K_0 K_0'' + K_0^{12} - K_0')(\ln x) + (K_0 K_0'' - K_0^{12} - 5/2K_0' + 2)(\ln x)^2 - \frac{1}{6}Q(\ln x)] \quad (30)$$

$$KK'' = 3K' + \frac{P}{k} - 3 - K^{12} + \frac{K_0}{xk} (K_0 K_0'' + K_0^{12} - 3K_0' + 3) \quad (31)$$

$$K^2 K''' = 3KK'' + \frac{(K - PK')}{K} - 3K'KK'' - \frac{K_0(K' - 1)}{xK} (K_0 K_0'' + K_0^{12} - 3K_0' + 3) \quad (32)$$

where $x = V/V_0$ and $Q = K_0 K_0'' + K_0^{12} - 3K_0' + 3$

Generalized Vinet-Rydberg Eos

Stacey [19, 20] has generalized the Vinet EOS, so as to make it compatible with infinite pressure value K'_∞ , for the pressure derivative of bulk modulus the equation thus formulated by Stacey is known is the generalized Rydberg EOS.

$$P = 3K_0 x^{-k_\infty^{-1}} (1 - x^{1/3}) \exp[\eta(1 - x^{1/3})] \quad (33)$$

$$K = 3K_0 x^{-k_\infty^{-1}} \exp[\eta(1 - x^{1/3})][k_\infty^{-1}(1 - x^{1/3}) + \frac{1}{3}x^{1/3} + \frac{\eta}{3}x^{1/3}(1 - x^{1/3})] \quad (34)$$

$$K' = \frac{3K_0 x^{-K'_\infty}}{K} \exp[\eta - x^{1/3}] [K_\infty^{12}(1-x^{1/3}) + x^{1/3} \left(\frac{1+\eta}{3}\right) \left(2K'_\infty - \frac{1}{3}\right) - x^{2/3} \eta (2K'_\infty - 1) + \frac{\eta^2}{9} x^{2/3} (1-x^{1/3})] \quad (35)$$

$$K'' = \frac{P}{K} \left[\frac{n}{27} x^{1/3} + \frac{q}{9} + q^2 + 2q^3 \right] + 2 \left(\frac{K}{P} \right) \left(K' - \frac{K}{P} \right) - K' \left(K' - \frac{K}{P} \right) \quad (36)$$

$$K''' = \frac{P}{K} \left[\frac{-n}{81} x^{1/3} + \frac{q}{27} - \frac{7}{9} q^2 + 4q^3 - 6q^4 \right] - 4KK'' \left(K' - \frac{K}{P} \right) - K^{12} \left(K' - \frac{K}{P} \right) + 6K' \left(\frac{K}{P} \right) \left(K' - \frac{K}{P} \right) - 6 \left(\frac{K}{P} \right)^2 \left(K' - \frac{K}{P} \right) \quad (37)$$

where $x = V/V_0$ and $\eta = \frac{3}{2} K'_0 - 3K'_\infty + \frac{1}{2} = -3K_0$

$$K''_0 - \frac{3}{4} K_0^{12} + \frac{1}{12}$$

$$q = \frac{x^{1/3}}{3(1-x^{1/3})}, \text{ for } K'_\infty = 2/3, \text{ Eq. (33) reduces}$$

to the original Rydberg EOS.

The Birch-Murnaghan EOS, the logarithmic EOS and the generalized Rydberg EOS can be written in the following form

$$K/P = K'_\infty + F(x) \quad (38)$$

Differentiating Eq (38) with the respect to 'P' successfully

$$(K' - K/P)K/P = -xF'(x) \quad (39)$$

$$[(KK'' + K'^2)(K/P) - 3K'(K/P)^2 + 2(K/P)^3 = xF'(x) + x^2F''(x)] \quad (40)$$

$$[(K^2K'''(K/P) + 4KK''(K' - K/P)(K/P) + K'^2(K' - K/P)(K/P) - 6K'(K' - K/P)(K/P)^2 + 6(K' - K/P)(K/P)^3] = -[x F'(x) + 3x^2F''(x) + x^3F'''(x)] \quad (41)$$

Where $x = V/V_0$

In case of birch murnagham fourth order EOS the value of F(x) is

$$F(x) = \left[\frac{2Ax^2 - \left(\frac{4}{3}\right)Bx^{\frac{4}{3}} + \left(\frac{2}{3}\right)Cx^{\frac{2}{3}}}{Ax^2 - Bx^{\frac{4}{3}} + Cx^{\frac{2}{3}} - D} \right] \quad (42)$$

In case of the logarithmic fourth -order EOS the value of F(x) is

$$F(x) = \left[\frac{1 - 2A_1 \ln x + 3A_2 (\ln x)^2}{-\ln x + A_1 (\ln x)^2 + A_2 (\ln x)^3} \right] \quad (43)$$

In case of the generalized Rydberg EOS the value of F(x) is

$$F(x) = \frac{\frac{1}{x^3}}{3(1-x^{\frac{1}{3}})} + \eta \frac{x^{\frac{1}{3}}}{3} \quad (44)$$

Results and Discussions

We can derive expressions for the derivatives of F(x) such as F'(x), F''(x) and F'''(x) by using Eqs. (42-44). In the limit $V \rightarrow 0$, $P \rightarrow \infty$, $K \rightarrow \infty$, but their ratio P/K remain finite such that $(P/K)_\infty = 1/K'_\infty$. Also $(1 - K'P/K)$, KK'' and K^2K''' tend to zero in the limit of infinite pressure, but their ratios $KK''(1 - K'P/K)$ and K^2K'''/KK'' remain finite [6,22]. At extreme compression $x \rightarrow 0$, we have $F(x) \rightarrow 0$, $x F'(x) \rightarrow 0$, $x^2 F''(x) \rightarrow 0$ and $x^3 F'''(x) \rightarrow 0$ for all the equations of state based on Eqs. (38-41) using the calculus, we have

$$\frac{xF'(x) + x^2F''(x)}{xF'(x)} = \frac{F'(x) + xF''(x) + 2xF'''(x) + x^2F''''(x)}{F'(x) + xF''(x)} \quad (45)$$

In the extreme compression limit Eqs. (39) and (40) gives

$$\left[\frac{KK''}{K' - \frac{K}{P}} \right]_{\infty} - K'_\infty = - \frac{xF'(x) + x^2F''(x)}{xF'(x)} \quad (46)$$

Eqs. (40) and (41) gives

$$\frac{\left(\frac{K^2 K'''}{KK''}\right)_{\infty} + K'_{\infty} \left(\frac{K' - K/P}{KK''}\right)_{\infty}}{1 - K'_{\infty} \left(\frac{K' - K/P}{KK''}\right)_{\infty}} = \frac{x F'(x) + 3x^2 F''(x) + x^3 F'''(x)}{x F'(x) + x^2 F''(x)} \quad (47)$$

Eqs. (45 – 47) then yield

$$\left(\frac{K^2 K'''}{KK''}\right)_{\infty} = \left(\frac{KK''}{K' - K/P}\right)_{\infty} - 2K'_{\infty} \quad (48)$$

The Birch-Murnaghan Fourth Order EOS gives using Eqs. (19-23)

$$\left(\frac{KK''}{K' - K/P}\right)_{\infty} = -K'_{\infty}(K'_{\infty} - 2/3) \quad (49)$$

$$\left(\frac{K^2 K'''}{KK''}\right)_{\infty} = -(K'_{\infty} + 2/3) \quad (50)$$

The logarithmic fourth order EOS using Eqs. (28-32)

$$\left(\frac{KK''}{K' - K/P}\right)_{\infty} = -2K'_{\infty} \quad (51)$$

$$\left(\frac{K^2 K'''}{KK''}\right)_{\infty} = -K'_{\infty} \quad (52)$$

The generalized Vinet-Rydberg EOS using Eqs. (33-37) gives

$$\left(\frac{KK''}{K' - K/P}\right)_{\infty} = \left(K'_{\infty} - \frac{1}{3}\right) \quad (53)$$

$$\left(\frac{K^2 K'''}{KK''}\right)_{\infty} = \left(K'_{\infty} + \frac{1}{3}\right) \quad (54)$$

All these equations of states satisfies the common relation (48). This relationship can be useful for investigating further the thermoelastic properties of solids at high pressures [6, 8, 23, 24].

References

1. Anderson, O.L.; Masuda, K.; Isaak, D.G. "A new thermodynamic approach for high-pressure physics." *Physics of the Earth and Planetary Interiors* 91.1-3 (1995): 3-16.
2. Shanker, J.; Thomas, L.M. *Phys. Stat. Sol (b)* 35 (1997): 601.
3. Chauhan, R. S., Singh, C. P. "Equation of state and thermal expansivity of NaCl under high pressure and high temperature." *Physica B: Condensed Matter* 387.1-2 (2007): 352-357.
4. Shanker, J ; Singh, P K; Saurabh, S. "Analysis of infinite pressure behaviour of thermoelastic properties of materials" *Indian Journal of Pure and Applied Physics*. 539(4) (2015): 230-233.
5. Ahmad, J.F.; Alkammash, I.Y. "Theoretical study of some thermodynamical properties for solid under high pressure using finite-strain EOS." *Journal of the Association of Arab Universities for Basic and Applied Sciences* 12.1 (2012): 17-22.
6. Shanker, J., Singh, B.P. "A comparative study of Keane's and Stacey's equations of state." *Physica B: Condensed Matter* 370.1-4 (2005): 78-83.
7. Kushwah, S.S., Bhardwaj, N.K. "Analysis based on equation of state for sodium halides." *Journal of Physics and Chemistry of Solids* 70.3-4 (2009): 700-702.
8. Stacey, F.D., Davis, P.M. "High pressure equations of state with applications to the lower mantle and core." *Physics of the Earth and Planetary interiors* 142.3-4 (2004): 137-184.
9. Stacey, F.D. "High pressure equations of state and planetary interiors." *Reports on progress in physics* 68.2 (2005): 341.
10. Vocadlo, L.; Poirer, J.P.; Price, G.D. "Grüneisen parameters and isothermal equations of state." *American Mineralogist* 85.2 (2000): 390-395.
11. Zubarev, V.N.; Vashchenko, V.Ya. "O koefficiente Gryunajzena" *Soviet Phys. Solid State* 5 (1963): 653. (in Russian).
12. J.C. Slater, *Introduction to chemical Physics*, Mc Graw – Hills, New York (1939).
13. Dugdale, J.S., MacDonald, D.K.C. "The thermal expansion of solids." *Physical Review* 89.4 (1953): 832.
14. Barton, M.A., Stacey, F.D. "The Grüneisen parameter at high pressure: a molecular dynamical study." *Physics of the earth and planetary interiors* 39.3 (1985): 167-177.

15. Holzapfel, W.B., Hartwig, M.; Sievers, W. "Equations of state for Cu, Ag, and Au for wide ranges in temperature and pressure up to 500 GPa and above." *Journal of Physical and Chemical Reference Data* 30.2 (2001): 515-529.
16. Anderson O.L. *Equations of State of Solid for Geophysics and Ceramic Science*, Oxford University Press, New York (1995).
17. Birch, F. "Elasticity and constitution of the Earth's interior." *Journal of Geophysical Research* 57 (1952): 227-286.
18. Poirier, J.P.; Tarantola, A. "A logarithmic equation of state" *Physics of the Earth and Planetary Interiors* 109 (1998).
19. Keane, A. "An investigation of finite strain in an isotropic material subjected to hydrostatic pressure and its seismological applications." *Australian Journal of Physics* 7.2 (1954): 322-333.
20. Stacey, F.D. "Finite strain, thermodynamics and the earth's core." *Physics of the Earth and Planetary Interiors* 128.1-4 (2001): 179-193.
21. Stacey, F.D. "The K-primed approach to high-pressure equations of state." *Geophysical Journal International* 143.3 (2000): 621-628.
22. Shanker, J.; Singh, B.P.; Jitendra, K. "Extreme compression behaviour of higher derivative properties of solids based on the generalized Rydberg equation of state." *Condense Matter Phys.* 12 (2009): 205-213.
23. Shanker, J.; Dulari, P.; Singh, P.K. "Extreme compression behaviour of equations of state." *Physica B: Condensed Matter* 404.21 (2009): 4083-4085.
24. K.S. Singh. "Second order and third order Grüneisen parameters at extreme compression." *Physica B: Condensed Matter* 407 (2012): 668-669